Study Title

Glutamic Acid and GABA - Occurrence of Decarboxylation of Glutamate to GABA

Literature Citation: Biological Chemistry, H. R. Mahler and E. H. Cordes Harper & Row Publishers, Copyright 1966, Page 685

Data Requirement

Not Applicable

Authors

H. R. Mahler and E. H. Cordes

Study Completed on


Not Applicable

Performing Laboratory

Not Applicable

Laboratory Project ID Number

Not Applicable



Statement of Data Confidentiality

No claim of confidentiality is made for any information contained in this study on the basis of its falling within the scope of FIFRA Section 10(d)(1)(A), (B) or (C).

Company:

Auxein Corporation

3900 Collins Road P. O. Box 27519

Lansing, Michigan 48909-0519

Company Agent:

M. Olav Messerschmidt

Vice-President, Business Development

Signature:

M. alder Missesschmidh

Date: 17, 1997

Good Laboratory Practices Statement

The submitter of this study was neither the sponsor of this study nor conducted it, and does not know whether it has been conducted in accordance with 40 CFR Part 160.

Submitter:

Auxein Corporation

3900 Collins Road P. O. Box 27519

Lansing, Michigan 48909-0519

Company Agent:

M. Olav Messerschmidt

Vice-President, Business Development

BIOLOGICAL CHEMISTRY

HENRY R. MAHLER & EUGENE H. CORDES DEPARTMENT OF CHEMISTRY, INDIANA UNIVERSITY

HARPER & ROW, PUBLISHERS, NEW YORK AND LONDON

To Annemarie and Shirley

BIOLOGICAL CHEMISTRY
by Henry R. Mahler and Eugene H. Cordes
Copyright @ 1966

Printed in the United States of America. All rights reserved. No part of this book may be used or reproduced in any manner whatsoever without written permission except in the case of brief quotations embodied in critical articles and reviews.

For information address Harper & Row, Publishers, Incorporated, 49 East 33rd Street, New York, N. Y. 10016.

Library of Congress Catalog Card Number: 66-19243

The metabolic roles of transaminations are several and significant. Important functions include (1) amino acid biosynthesis (transamination is directly involved in the synthesis of at least eleven of the amino acids); (2) amino acid degradation (see discussion below); (3) liaison between carbohydrate and amino acid metabolism; and (4) synthesis of several specific compounds, including urea and γ -amino butyric acid.

The transaminases are ubiquitous enzymes. Individual enzymes may be fairly specific for a given substrate pair or, alternatively, may be able to catalyze reactions involving a broad spectrum of amino and keto acids. At

TABLE 16-5. Enzymic Decarboxylation of Amino Acids

Amino Acid	Decarboxylation produ	
L-Arginine L-Aspartic acid L-Cysteic acid L-Cysteine sulfinic acid L-Glutamic acid	Agmatine β-Alanine Taurine Hypotaurine γ-Aminobutyric acid	Microorganisms (e.g., E. coli Rhizobium leguminosarum Liver, spleen, brain Liver, spleen, brain Microorganisms (e.g., Cl. welchii F. coli)
γ-Hydroxy-L-glutamic acid	acid d-nyuroxyour	plants (barley spinesh attack
γ-Methylene-L-glutamic ac	acid	tyric Barley, red pepper, peanut
L-Lysine	Histamine Cadaverine	Microorganisms (e.g., Cl. welchit, Lactobactlit), animal tissues (kidney, liver, duodenum) Microorganisms (e.g., B.
meso-α, ε-Diaminopimelic acid	L-Lysine	Microorganisms (e.g. F. a.)
L-Ornithine	Putrescine	Microorganisms (e.g. Cl
t-Phenylalanine	eta-Phenylethylamine	Microorganisms (a a stand
L-Tyrosine 3,4-Dihydroxy-L-phenyl-	Tyramine	tococcus fecalis) Microorganisms (e.g., S. fecalis), animal tissues (e.g., kidney)
-Hydroxy-1-tryptophan	3,4-Dihydroxy-β-phenyl- ethylamine 5-Hydroxyfryptamine	Microorganisms (e.g., S. fecalis), animal tissues (e.g., kidney)
Valine	Isobutylamine	Animal tissues (e.g., kidney, brain, stomach)
Leucine Lysine	Isoamylamine	Proteus vulgaris Proteus vulgaris
URCE: J. Fruton and S. Simon	ids: "General Diagram	Microorganisms